#: B-040a
Page: 1 of 1
Orig. Date: 10/2008
Revised: 09/11/2019

INDICATION:

Pulseless cardiopulmonary arrest Adolescent/Adult to provide continuous chest compression.

CONTRAINDICATIONS:

- Chest circumference beyond the fit range for the compression band.
- Recent chest surgery or a chest tube in place.
- Left ventricular assist device in place.

PROCEDURE:

- Remove clothing from thorax area.
- Apply defibrillation-monitor pads to chest in usual location and manner. If device is pass through defibrillator capable, follow manufacturer instructions for specific device.
- Log roll technique:
 - o Extend and open device next to patient
 - o Log roll patient away from device
 - O Slide device under back or patient and roll onto support area of device band
 - o Aline and close compression band fully extended on patient chest.
 - o Activate device and assess for proper functioning.
- Tilt forward technique:
 - o Extend and open device next to patient
 - o Lean (sit) patient forward
 - o Slide device behind patient and position down to buttocks
 - o Position device, check alignment and close compression band over patient's chest
 - Activate device and assess for proper functioning.

DOCUMENTATION:

- Document time device placed.
- Check "AutoPulse® box on PCR.

Approved:

Review Date: 10/2008; 5/2018; 9/2019

Final Date for Implementation: 10/01/2019

OCEMS copyright © 2019